



## Human Cancer/Testis Antigen 1B (CTAG1B) Protein

Catalogue No.: abx166945



SDS-PAGE analysis of recombinant Human Cancer/Testis Antigen 1B Protein.



Gene sequencing analysis of recombinant Human Cancer/Testis Antigen 1B Protein.

Human Cancer/Testis Antigen 1B Protein is a recombinant Human protein expressed in *E. coli*.

**Target:** Cancer/Testis Antigen 1B (CTAG1B)

**Origin:** Human

**Expression:** Recombinant

**Tested Applications:** WB, SDS-PAGE

**Host:** *E. coli*

**Conjugation:** Unconjugated

**Form:** Lyophilized

**Purity:** > 90%

**Reconstitution:** To keep the original salt concentration, we recommend reconstituting to the original concentration prior to lyophilization (see Concentration) in ddH<sub>2</sub>O. If a lower concentration is required, dilute in 20 mM Tris, 150 mM NaCl, pH 8.0. If a higher concentration is required, the product can be reconstituted directly in 20 mM Tris, 150 mM NaCl, pH 8.0, though please note that this will change the overall salt concentration. The stock concentration should be between 0.1-1.0 mg/ml. Do not vortex.

# Datasheet

Version: 6.0.0  
Revision date: 30 Mar 2025



<b>Storage:</b>	Store at 2-8 °C for up to one month. Store at -80 °C for up to one year. Avoid repeated freeze/thaw cycles.
<b>Molecular Weight:</b>	Calculated MW: 21.7 kDa Observed MW (SDS-PAGE): 25 kDa Possible reasons why the actual band size differs from the predicted band size: <ol style="list-style-type: none"><li>1. Splice variants. Alternative splicing may create different sized proteins from the same gene.</li><li>2. Relative charge. The composition of amino acids may affect the charge of the protein.</li><li>3. Post-translational modification. Phosphorylation, glycosylation, methylation etc. may affect the band size.</li><li>4. Post-translational cleavage. Many proteins are synthesised as pro-proteins, and then cleaved to give the active form.</li><li>5. Polymerisation of the target protein. Dimerisation, multimerisation etc. will increase the band size observed.</li></ol>
<b>Sequence Fragment:</b>	Met1-Arg180
<b>Sequence:</b>	MQAEGRGTGG STGDADGPGG PGIPDGPGGN AGGPGEAGAT GGRGPRGAGA ARASGPGGGA PRGPHGGAA S GLNGCCRCGA RGPESRLLEF YLAMPFATPM EAELARRSLA QDAPPLPVPV VLLKEFTVSG NILTIRL TAA DHRQLQLSIS SCLQQLSLLM WITQCFLPVF LAQPPSGQRR
<b>Tag:</b>	N-terminal His tag
<b>Buffer:</b>	Prior to lyophilization: 20 mM Tris, 150 mM NaCl, pH 8.0, containing 1 mM EDTA, 0.01% Sarcosyl, 5% Trehalose.
<b>Activity:</b>	Not tested
<b>Concentration:</b>	Prior to lyophilization: 200 µg/ml
<b>Note:</b>	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.