



Fc Fragment Of IgA Receptor / CD89 (FCAR) Antibody


Catalogue No.: abx130285




Western blot analysis of Rat Serum.



Western blot analysis of recombinant Rat FcaR.



IHC-P analysis of Rat Kidney Tissue, with DAB staining.



IHC-P analysis of Rat Kidney Tissue, with DAB staining.

Fc Fragment Of IgA Receptor Antibody is a Rabbit Polyclonal against Fc Fragment Of IgA Receptor.

Target: Fc Fragment Of IgA Receptor / CD89 (FCAR)

Clonality: Polyclonal

Reactivity: Rat

Datasheet

Version: 4.0.0
Revision date: 12 Mar 2025



Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: FcaR (Thr51-Arg280)

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Buffer: 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only