



Neural Cell Adhesion Molecule 2 (NCAM2) Antibody


Catalogue No.: abx128183




Western blot analysis of recombinant Human NCAM2.



Western blot analysis of (1) Porcine Brain Tissue and (2) Rat Brain Tissue.



IHC-P analysis of Human Kidney Tissue, with DAB staining, using Rabbit Anti-Human NCAM2 Antibody (30 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).



IHC-P analysis of Human Cerebrum Tissue, with DAB staining, using Rabbit Anti-Human NCAM2 Antibody (30 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody ([abx400043](#), 2 µg/ml).

Neural Cell Adhesion Molecule 2 Antibody is a Rabbit Polyclonal against Neural Cell Adhesion Molecule 2.

Target: Neural Cell Adhesion Molecule 2 (NCAM2)

Clonality: Polyclonal

Reactivity: Human

Datasheet

Version: 2.0.0
Revision date: 11 Feb 2025



Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.5-3 µg/ml, IHC: 5-30 µg/ml, IF/ICC: 5-30 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx166894 - Recombinant NCAM2 (Leu21-Gln297) expressed in E. coli
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	O15394 (UniProt , ExPASy)
Gene Symbol:	NCAM2
GeneID:	4685
KEGG:	hsa:4685
String:	9606.ENSP00000383392
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Specificity:	Cross-reacts with Mouse, Rat and Pig NCAM2.
Concentration:	0.38 mg/ml
Note:	This product is for research use only.