



Nucleolin (NCL) Antibody

Catalogue No.: abx006806



Western blot analysis of extracts of various cell lines using NCL Antibody (1/1000 dilution).



Immunohistochemistry of paraffin-embedded human liver cancer using NCL Antibody (1/100 dilution, 40x lens). Microwave antigen retrieval was performed in 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.

NCL Antibody is a Rabbit Polyclonal Antibody against NCL.

Target: Nucleolin (NCL)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC-P: 1/50 - 1/200. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: Recombinant fusion protein corresponding to human NCL

Isotype: IgG

Form: Liquid

Purification: Purified by affinity chromatography.

Datasheet

Version: 4.0.0
Revision date: 30 Apr 2025



Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: P19338 ([UniProt](#), [ExPASy](#))

Gene Symbol: NCL

GeneID: [4691](#)

NCBI Accession: NP_005372.2

KEGG: hsa:4691

String: [9606.ENSP00000318195](#)

Molecular Weight: Calculated MW: 76 kDa
Observed MW: 110 kDa

Buffer: PBS, pH 7.3, containing 0.01% thiomersal, 50% glycerol.

Concentration: 1 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only