


Low Affinity Immunoglobulin Epsilon Fc Receptor / CD23 (FCER2) Antibody

Catalogue No.: abx001520



Western blot analysis of lysates from MCF-7 cells, using CD23 Antibody at 1/1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) at 1/10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

FCER2 Antibody is a Rabbit Polyclonal antibody against FCER2. The protein encoded by this gene is a B-cell specific antigen, and a low-affinity receptor for IgE. It has essential roles in B cell growth and differentiation, and the regulation of IgE production. This protein also exists as a soluble secreted form, then functioning as a potent mitogenic growth factor. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2011].

Target:	Low Affinity Immunoglobulin Epsilon Fc Receptor / CD23 (FCER2)
Clonality:	Polyclonal
Reactivity:	Human, Mouse
Tested Applications:	ELISA, WB
Host:	Rabbit
Recommended dilutions:	ELISA: 1 µg/ml, WB: 1/500 - 1/2000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 48-321 of human CD23.
Isotype:	IgG
Form:	Liquid
Purification:	Purified by affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P06734 (UniProt , ExpASY)

Datasheet

Version: 3.0.0
Revision date: 03 Feb 2025



Gene Symbol: FCER2

GeneID: [2208](#)

NCBI Accession: NP_001993.2

Molecular Weight: Calculated MW: 36 kDa
Observed MW: 36 kDa

Buffer: PBS, pH 7.3, containing 0.02% sodium azide, 50% glycerol.

Concentration: > 0.2 mg/ml

Note: This product is for research use only.

For Reference Only