



Somatostatin Receptor Type 2 (SSTR2) Antibody


Catalogue No.: abx317864




Western blot analysis of Mouse spleen tissue using SSTR2 Antibody (2.7 µg/ml) and Goat Polyclonal Anti-Rabbit IgG (1/50000 dilution). Calculated MW: 42 kDa, 41 kDa. Observed MW: 42 kDa.



IHC-P analysis of Human kidney tissue using SSTR2 Antibody (1/100 dilution).



IF analysis of HepG2 cells using SSTR2 Antibody (1/100 dilution) and AF488-conjugated Goat Anti-Rabbit IgG.



IF analysis of PC-3 cells using SSTR2 Antibody (1/100 dilution) and AF488-conjugated Goat Anti-Rabbit IgG.

SSTR2 Antibody is a Rabbit Polyclonal against SSTR2.

Target: Somatostatin Receptor Type 2 (SSTR2)

Clonality: Polyclonal

Reactivity: Human, Mouse

Datasheet

Version: 4.0.0
Revision date: 09 Apr 2025



Tested Applications:	ELISA, WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 1/1000 - 1/5000, IHC: 1/20 - 1/200, IF/ICC: 1/50 - 1/200. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant human Somatostatin receptor type 2 protein (304-369 AA).
Isotype:	IgG
Form:	Liquid
Purity:	> 95%
Purification:	Purified by Protein G.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P30874 (UniProt , ExPASy)
Gene Symbol:	SSTR2
GeneID:	6752
OMIM:	182452
NCBI Accession:	NP_001041.1
KEGG:	hsa:6752
String:	9606.ENSP00000350198
Molecular Weight:	Calculated MW: 42 kDa, 41 kDa Observed MW: 42 kDa
Buffer:	0.01 M PBS, pH 7.4, containing 0.03% Proclin-300 and 50% Glycerol.
Concentration:	4 mg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.