



T-Lymphocyte Activation Antigen CD86 / B7-2 (CD86) Antibody

Catalogue No.: abx231504



WB analysis of Jurkat cells, using CD86 antibody (1/500 dilution).



IHC-P analysis of human tonsillitis tissue, using CD86 Antibody (1/200 dilution). Heat-mediated antigen retrieval was carried out with Tris-EDTA buffer, pH 9.

CD86 Antibody is a Rabbit Polyclonal against CD86.

Target:	T-Lymphocyte Activation Antigen CD86 / B7-2 (CD86)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	ELISA, WB, IHC
Host:	Rabbit
Recommended dilutions:	WB: 1/200 - 1/2000, IHC: 1/50 - 1/500. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	CD86 molecule
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)

Datasheet

Version: 1.0.0
Revision date: 22 Nov 2024



Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
UniProt Primary AC:	P42081 (UniProt , ExpASY)
Gene Symbol:	CD86
KEGG:	hsa:942
String:	9606.ENSP00000332049
Molecular Weight:	Observed MW: 75 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	This product is for research use only.

For Reference Only