



CFLAR / FLIP Antibody


Catalogue No.: abx231621



WB analysis of Jurkat cells, using CFLAR antibody (1/1000 dilution).



IHC-P analysis of human skeletal muscle tissue, using CFLAR Antibody (1/50 dilution).



IP analysis of Jurkat cell lysate (4000 µg), using CFLAR/FLIP antibody (4 µg, detection: 1/500 dilution).

CFLAR/FLIP Antibody is a Rabbit Polyclonal against CFLAR/FLIP.

Target: CFLAR/FLIP

Clonality: Polyclonal

Reactivity: Human

Tested Applications: ELISA, WB, IHC, IF/ICC, IP

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/20 - 1/200, IF/ICC: 1/20 - 1/200, IP: 1/200 - 1/1000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Datasheet

Version: 1.0.0
Revision date: 27 Nov 2024



Immunogen:	CASP8 and FADD-like apoptosis regulator
Isotype:	IgG
Form:	Liquid
Purity:	≥ 95% (SDS-PAGE)
Purification:	Purified by immunogen affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Validity:	12 months.
Molecular Weight:	Observed MW: 30-35 kDa, 50-55 kDa
Buffer:	PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.
Concentration:	2 mg/ml
Note:	This product is for research use only.

For Reference Only