



G-Protein Coupled Receptor 94 (GPR94) Antibody


Catalogue No.: abx430396




abx430396 (0.03 µg/ml) staining of NIH3T3 lysate (35 µg protein in RIPA buffer). Detected by chemiluminescence.




abx430396 (0.1 µg/ml) staining of U2OS (A), HEK293 (B), HeLa (C) and (0.3 µg/ml) A549 cell lysate (35 µg protein in RIPA buffer). Detected by chemiluminescence.



abx430396 (5 µg/ml) staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer Ph 6, AP-staining.




abx430396 (3.8 µg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.




Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml), showing Endoplasmic reticulum and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).

Datasheet

Version: 2.0.0
Revision date: 28 Dec 2024



Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml), showing Endoplasmic reticulum and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 µg/ml) followed by AF488 secondary antibody (2 µg/ml).



Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 µg/ml) followed by AF488 secondary antibody (1 µg/ml). IgG control: Unimmunized goat IgG (black line) followed by AF488 secondary antibody.

GPR94 Antibody is a Goat Polyclonal antibody against GPR94.

Target:	G-Protein Coupled Receptor 94 (GPR94)
Clonality:	Polyclonal
Reactivity:	Human, Mouse
Tested Applications:	P-ELISA, WB, IHC, IF/ICC, FCM
Host:	Goat
Recommended dilutions:	P-ELISA: 1/32000. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx269926 - Internal region: C-KEGVKFDSEKTKE
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Datasheet

Version: 2.0.0
Revision date: 28 Dec 2024



Gene Symbol: HSP90B1

GeneID: [7184](#) [22027](#) [362862](#)

NCBI Accession: NP_003290.1

Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: This product is for research use only.

For Reference Only