



Allograft Inflammatory Factor 1 (AIF1) Antibody


Catalogue No.: abx430051



Western blot analysis of Human lymph node and Mouse brain lysates (35 µg protein in RIPA buffer) using Allograft Inflammatory Factor 1 Antibody (1 µg/ml).




Immunohistochemistry analysis of paraffin-embedded Human spleen tissue, using AIF1 antibody (6 µg/ml) and HRP staining. Heat induced antigen retrieval was carried out in citrate buffer, pH 6, prior to staining.



Negative Control showing staining of paraffin embedded Human spleen, with no primary antibody.



Immunohistochemistry analysis of paraffin-embedded Rat spleen tissue, using AIF1 antibody (8 µg/ml) and HRP staining. Heat induced antigen retrieval was carried out in citrate buffer, pH 6, prior to staining.



Negative Control showing staining of paraffin embedded Rat spleen, with no primary antibody.

Datasheet

Version: 2.0.0
Revision date: 06 Oct 2024



Allograft Inflammatory Factor 1 (AIF1) (Isoform 1 and 3) Antibody is a Goat Polyclonal antibody against Allograft Inflammatory Factor 1 (AIF1).

Target:	Allograft Inflammatory Factor 1 (AIF1)
Clonality:	Polyclonal
Reactivity:	Human, Mouse, Rat
Tested Applications:	P-ELISA, WB, IHC
Host:	Goat
Recommended dilutions:	P-ELISA: 1/128000, WB: 0.3-1 µg/ml, IHC: 4-6 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx269155 - C-Terminus: C-TGPPAKKAISELP
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	AIF1
GeneID:	199 11629 29427
NCBI Accession:	NP_116573.1, NP_001614.3
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.
Concentration:	0.5 mg/ml
Note:	This product is for research use only.