



Proliferating Cell Nuclear Antigen (PCNA) Antibody


Catalogue No.: abx431497




abx431497 (0.05 µg/ml) staining of Jurkat lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



abx431497 (0.03 µg/ml) staining of NIH3T3 (A), Mouse Testis (B) and Rat Testis (C) lysates (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



abx431497 (0.3 µg/ml) staining of Pig Spleen lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



abx431497 (2.5 µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

PCNA Antibody is a Goat Polyclonal antibody against PCNA.

Target: Proliferating Cell Nuclear Antigen (PCNA)

Clonality: Polyclonal

Reactivity: Human, Mouse, Rat, Pig

Datasheet

Version: 2.0.0
Revision date: 03 Jan 2025



Tested Applications:	P-ELISA, WB, IHC
Host:	Goat
Recommended dilutions:	P-ELISA: 1/128000, WB: 0.03-0.1 µg/ml, IHC: 2.5 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	abx616871 - Internal region, 111-122 AA: C-EAPNQEKVSDYE
Isotype:	IgG
Form:	Liquid
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
Gene Symbol:	PCNA
GeneID:	5111 18538 25737
NCBI Accession:	NP_002583.1
Buffer:	Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.
Concentration:	0.5 mg/ml
Note:	This product is for research use only.

For Reference Only