



Transcription Factor AP-2-Delta (TFAP2D) Antibody


Catalogue No.: abx431730




IF analysis of A549 cells, using Transcription Factor AP-2-Delta (TFAP2D) Antibody (10 ug/ml) followed by secondary antibody conjugated to Alexa Fluor 488 (2ug/ml). DAPI was used for nuclear staining. Non-specific Goat IgG antibody was used as a negative control (10 ug/ml).



IF analysis of U2OS cells, using Transcription Factor AP-2-Delta (TFAP2D) Antibody (10 ug/ml) followed by secondary antibody conjugated to Alexa Fluor 488 (2ug/ml). DAPI was used for nuclear staining. Non-specific Goat IgG antibody was used as a negative control (10 ug/ml).



WB analysis of NIH3T3 nuclear cell lysate, using Transcription Factor AP-2-Delta (TFAP2D) Antibody (1 µg/ml).



WB analysis of Human Ovary lysate, using Transcription Factor AP-2-Delta (TFAP2D) Antibody (2 µg/ml).

TFAP2D Antibody is a Goat Polyclonal antibody against TFAP2D.

Target: Transcription Factor AP-2-Delta (TFAP2D)

Clonality: Polyclonal

Tested Applications: P-ELISA, WB, IF/ICC, FCM

Datasheet

Version: 3.0.0
Revision date: 19 Mar 2025



Host: Goat

Recommended dilutions: P-ELISA: 1/64000. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: [abx615688](#) - Internal region: C-HHQSFHVEFQHSHP

Isotype: IgG

Form: Liquid

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Gene Symbol: TFAP2D

GeneID: [83741](#) [226896](#) [301284](#)

NCBI Accession: NP_758438.2

Buffer: Tris saline, pH 7.3, containing 0.02% sodium azide and 0.5% bovine serum albumin.

Concentration: 0.5 mg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.