



Noggin (NOG) Antibody


Catalogue No.: abx132077



Western blot analysis of Rat Placenta lysate, using Human NOG Antibody (2 µg/ml) and HRP-conjugated Goat Anti-Mouse antibody ([abx400001](#), 0.2 µg/ml).



Western blot analysis of recombinant Human NOG.



IHC-P analysis of Human Liver cancer Tissue, with DAB staining.



IHC-P analysis of Human Liver cancer Tissue, with DAB staining.

Noggin Antibody is a Mouse Monoclonal against Noggin.

Target: Noggin (NOG)

Clonality: Monoclonal

Reactivity: Human

Datasheet

Version: 1.0.0
Revision date: 06 Oct 2024



Tested Applications:	WB, IHC, IF/ICC
Host:	Mouse
Recommended dilutions:	WB: 0.5-5 µg/ml, IHC: 5-30 µg/ml, IF/ICC: 5-30 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Gln28-Cys232
Isotype:	IgG _{2b} Kappa
Form:	Liquid
Purification:	Purified by Protein A and Protein G affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q13253 (UniProt , ExPASy)
Gene Symbol:	NOG
GeneID:	9241
OMIM:	184460
HGNC:	7866
KEGG:	hsa:9241
Ensembl:	ENSG00000183691
String:	9606.ENSP00000328181
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Note:	This product is for research use only.