



Activin Receptor Type IA Antibody

Catalogue No.: abx033179



Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. ACVR1 is an activin A type I receptor which signals a particular transcriptional response in concert with activin type II receptors.

Target: Activin Receptor Type IA

Clonality: Polyclonal

Reactivity: Human, Rat

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/1000, IHC-P: 1/50 - 1/100. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.

Conjugation: Unconjugated

Immunogen: KLH-conjugated synthetic peptide between 6-34 amino acids from the N-terminal region of human Activin Receptor Type IA (ACVR1).

Datasheet

Version: 1.0.0
Revision date: 15 Jan 2025



Isotype:	IgG
Form:	Liquid
Purification:	Purified Rabbit Polyclonal Antibody.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q04771 (UniProt , ExpASY)
KEGG:	hsa:90
String:	9606.ENSP00000263640
Molecular Weight:	Calculated MW: 57.2 kDa
Buffer:	PBS containing 0.09% sodium azide.
Specificity:	Predicted to react with Cow ACVR1.
Note:	This product is for research use only.

For Reference Only