


Proteasome Subunit Alpha Type 7 (PSMA7) Antibody

Catalogue No.:abx034050





The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMA7 is a member of the peptidase T1A family, that is a 20S core alpha subunit. This particular subunit has been shown to interact specifically with the hepatitis B virus X protein, a protein critical to viral replication. In addition, this subunit is involved in regulating hepatitis virus C internal ribosome entry site (IRES) activity, an activity essential for viral replication. This core alpha subunit is also involved in regulating the hypoxia-inducible factor-1alpha, a transcription factor important for cellular responses to oxygen tension.

Target: Proteasome Subunit Alpha Type 7 (PSMA7)

Clonality: Polyclonal

Reactivity: Human, Mouse

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/1000, IHC-P: 1/10 - 1/50. Not tested in IHC-F. Optimal dilutions/concentrations should be

determined by the end user.

Conjugation: Unconjugated

Immunogen: KLH-conjugated synthetic peptide between 100-126 amino acids from the Central region of human

PSMA7.

1 of 2

Datasheet

Version: 3.0.0 Revision date: 22 Dec 2024



Isotype: IgG

Form: Liquid

Purification: Purified through a protein A column, followed by peptide affinity purification.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

UniProt Primary AC: O14818 (<u>UniProt</u>, <u>ExPASy</u>)

Gene Symbol: PSMA7

KEGG: hsa:5688

String: <u>9606.ENSP00000359910</u>

Molecular Weight: Calculated MW: 27.9 kDa

Buffer: PBS containing 0.09% sodium azide.

Specificity: Predicted to react with Rat, Cow and Xenopus PSMA7.

Note: This product is for research use only.