


CD209 Antigen (CD209) Antibody

Catalogue No.:abx034379



CD209 encodes a transmembrane receptor and is often referred to as DC-SIGN because of its expression on the surface of dendritic cells and macrophages. The encoded protein is involved in the innate immune system and recognizes numerous evolutionarily divergent pathogens ranging from parasites to viruses with a large impact on public health. The protein is organized into three distinct domains: an N-terminal transmembrane domain, a tandem-repeat neck domain and C-type lectin carbohydrate recognition domain. The extracellular region consisting of the C-type lectin and neck domains has a dual function as a pathogen recognition receptor and a cell adhesion receptor by binding carbohydrate ligands on the surface of microbes and endogenous cells. The neck region is important for homo-oligomerization which allows the receptor to bind multivalent ligands with high avidity. Variations in the number of 23 amino acid repeats in the neck domain of this protein are rare but have a significant impact on ligand binding ability. This gene is closely related in terms of both sequence and function to a neighboring gene (GeneID 10332; often referred to as L-SIGN). DC-SIGN and L-SIGN differ in their ligand-binding properties and distribution.

Target:	CD209 Antigen (CD209)	
Clonality:	Polyclonal	
Reactivity:	Human	
Tested Applications:	ELISA, WB	
Host:	Rabbit	
Recommended dilutions: WB: 1/1000. Optimal dilutions/concentrations should be determined by the end user.		
Conjugation:	Unconjugated	
Immunogen:	KLH-conjugated synthetic peptide between 330-355 amino acids from the Central region of human CD209.	
lsotype:	lgG	
Form:	Liquid	
Purification:	Purified through a protein A column, followed by peptide affinity purification.	

Datasheet Version: 3.0.0 Revision date: 12 Mar 2025



Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	Q9NNX6 (<u>UniProt</u> , <u>ExPASy</u>)
Gene Symbol:	CD209
String:	<u>9606.ENSP00000315477</u>
Molecular Weight:	Calculated MW: 45.8 kDa
Buffer:	PBS containing 0.09% sodium azide.
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.