

## DCC-Interacting Protein 13-Beta (APPL2) Antibody

Catalogue No.:abx036769

Rabbit Polyclonal against the APPL2 protein.

Target:	DCC-Interacting Protein 13-Beta (APPL2)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	ELISA, WB, IHC
Host:	Rabbit
Recommended dilutions	: ELISA: 1/20000 - 1/80000, WB: 1/500 - 1/2000, IHC: 1/100 - 1/200. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	Recombinant fragment corresponding to 104-324 AA of human APPL2.
lsotype:	lgG
Form:	Lyophilized
Purification:	
	Purified by antigen affinity column chromatography.
Reconstitution:	Purified by antigen affinity column chromatography. Reconstitute in 100 $\mu$ l of sterile distilled H <sub>2</sub> O with 50% glycerol.
Reconstitution:	Reconstitute in 100 $\mu I$ of sterile distilled $H_2O$ with 50% glycerol.
Reconstitution: Storage:	Reconstitute in 100 $\mu$ l of sterile distilled H <sub>2</sub> O with 50% glycerol. Store at -20 °C. Avoid repeated freeze/thaw cycles.
Reconstitution: Storage: Molecular Weight:	Reconstitute in 100 µl of sterile distilled H₂O with 50% glycerol. Store at -20 °C. Avoid repeated freeze/thaw cycles. Observed MW: 74 kDa
Reconstitution: Storage: Molecular Weight: Buffer:	Reconstitute in 100 µl of sterile distilled H <sub>2</sub> O with 50% glycerol. Store at -20 °C. Avoid repeated freeze/thaw cycles. Observed MW: 74 kDa Prior to lyophilization: 1% BSA and 0.02% NaN3. Predicted to react with Mouse and Rat APPL2. Lyophilized form: Not applicable.
Reconstitution: Storage: Molecular Weight: Buffer: Specificity:	Reconstitute in 100 µl of sterile distilled H₂O with 50% glycerol. Store at -20 °C. Avoid repeated freeze/thaw cycles. Observed MW: 74 kDa Prior to lyophilization: 1% BSA and 0.02% NaN3. Predicted to react with Mouse and Rat APPL2.