



Human Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL) Protein (Active)

Catalogue No.: abx651459



Human Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL) Protein (Active)



Gene sequence extract of Human Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL) Protein (Active).

Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL) Protein (Active) is an active protein from Human.

Target: Tumor Necrosis Factor Related Apoptosis Inducing Ligand (TRAIL)

Origin: Human

Tested Applications: WB, SDS-PAGE

Conjugation: Unconjugated

Form: Lyophilized

Purity: > 95%

Reconstitution: To keep the original salt concentration, we recommend reconstituting to the original concentration prior to lyophilization (see Concentration) in ddH₂O. If a lower concentration is required, dilute in 20 mM Tris, 150 mM NaCl, pH 8.0. If a higher concentration is required, the product can be reconstituted directly in 20 mM Tris, 150 mM NaCl, pH 8.0, though please note that this will change the overall salt concentration. The stock concentration should be between 0.1-1.0 mg/ml. Do not vortex.

Storage: Store at 2-8 °C for up to one month. Store at -80 °C for up to one year. Avoid repeated freeze/thaw cycles.

Datasheet

Version: 3.0.0

Revision date: 07 Apr 2025



UniProt Primary AC: Q6IBA9 ([UniProt](#), [ExpASY](#))

Molecular Weight: Calculated MW: 19.6 kDa
Observed MW: 19 kDa

Sequence Fragment: Val114-Gly281

Tag: N-terminal His tag

Buffer: Prior to lyophilization: 20 mM Tris, 150 mM NaCl, pH 8.0, containing 1 mM EDTA, 1 mM DTT, 0.01% Sarcosyl, 5% Trehalose and Proclin-300.

Activity: Active

Concentration: Prior to lyophilization: 200 µg/ml

Note: THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only