


Human Microfibrillar Associated Protein 2 (MFAP2) Protein (Active)

Catalogue No.:abx651415



Human Microfibrillar Associated Protein 2 (MFAP2) Protein (Active)



Gene sequence extract of Human Microfibrillar Associated Protein 2 (MFAP2) Protein (Active).

Microfibrillar Associated Protein 2 (MFAP2) Protein (Active) is an active protein from Human.

Target:	Microfibrillar Associated Protein 2 (MFAP2)
Origin:	Human
Tested Applications	: WB, SDS-PAGE
Host:	E. coli
Conjugation:	Unconjugated
Form:	Lyophilized
Purity:	> 95%
Reconstitution:	To keep the original salt concentration, we recommend reconstituting to the original concentration prior to lyophilization (see Concentration) in ddH_2O . If a lower concentration is required, dilute in 20 mM Tris, 150 mM NaCl, pH 8.0. If a higher concentration is required, the product can be reconstituted directly in 20 mM Tris, 150 mM NaCl, pH 8.0, though please note that this will change the overall salt concentration. The stock concentration should be between 0.1-1.0 mg/ml. Do not vortex.
Storage:	Store at 2-8 °C for up to one month. Store at -80 °C for up to one year. Avoid repeated freeze/thaw cycles.



UniProt Primary AC: P55001 (UniProt, ExPASy)

KEGG:	hsa:4237
String:	<u>9606.ENSP00000364685</u>
Molecular Weight:	Calculated MW: 22.0 kDa Observed MW: 38 kDa
Sequence Fragment	: Leu6-Val162
Tag:	N-terminal His tag
Buffer:	Prior to Iyophilization: 20 mM Tris, 150 mM NaCl, pH 8.0, containing 1 mM EDTA, 1 mM DTT, 0.01% Sarcosyl, 5% Trehalose and Proclin-300.
Activity:	Active
Concentration:	Prior to lyophilization: 200 µg/ml
Note:	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.