

## Fc Fragment Of IgA Receptor / CD89 (FCAR) Antibody (FITC)

Catalogue No.: abx270386



Fluorescence emission spectra of FITC.

Fc Fragment Of IgA Receptor (FcaR) Antibody (FITC) is a Rabbit Polyclonal Antibody conjugated to FITC against Fc Fragment Of IgA Receptor (FcaR) for use in flow cytometry.

|                               |  |
|-------------------------------|--|
| <b>Target:</b>                | Fc Fragment Of IgA Receptor / CD89 (FCAR)  |
| <b>Clonality:</b>             | Polyclonal   |
| <b>Reactivity:</b>            | Human  |
| <b>Tested Applications:</b>   | FCM  |
| <b>Host:</b>                  | Rabbit   |
| <b>Recommended dilutions:</b> | FCM: 1-5 $\mu$ l/10 <sup>6</sup> cells. Optimal dilutions/concentrations should be determined by the end user. |
| <b>Conjugation:</b>           | FITC   |
| <b>Excitation/Emission:</b>   | 499/515  |
| <b>Laser Line:</b>            | 488  |
| <b>Isotype:</b>               | IgG  |
| <b>Form:</b>                  | Liquid   |
| <b>Purification:</b>          | Purified by antigen-specific affinity chromatography.  |
| <b>Storage:</b>               | Aliquot and store at -20°C. Avoid exposure to light. Avoid repeated freeze/thaw cycles.                        |
| <b>Buffer:</b>                | 0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.  |

# Datasheet

Version: 2.0.0

Revision date: 12 Mar 2025



**Note:**

THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only