


Alpha-2-Heremans Schmid Glycoprotein (AHSG) Antibody


Catalogue No.:abx175340




Western blot analysis of (1) Cow Serum, (2) Cow Liver lysate, and (3) Human HepG2 cell lysate.




IHC-P analysis of Cow Cerebrum Tissue, with DAB staining, using Rabbit Anti-Cow AHSG Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody (abx400043, 2 µg/ml).



IHC-P analysis of Cow Liver Tissue, with DAB staining, using Rabbit Anti-Cow AHSG Antibody (20 μg/ml) and HRP-conjugated Goat Anti-Rabbit antibody (abx400043, 2 μg/ml).



IHC-P analysis of Cow Kidney Tissue, with DAB staining, using Rabbit Anti-Cow AHSG Antibody (20 μg/ml) and HRP-conjugated Goat Anti-Rabbit antibody (abx400043, 2 μg/ml).



IHC-P analysis of Cow Cerebellum Tissue, with DAB staining, using Rabbit Anti-Cow AHSG Antibody (20 µg/ml) and HRP-conjugated Goat Anti-Rabbit antibody (abx400043, 2 µg/ml).

Datasheet

Version: 3.0.0 Revision date: 03 Dec 2024



Alpha-2-Heremans Schmid Glycoprotein (AHSG) Antibody is a Rabbit Polyclonal antibody against Alpha-2-Heremans Schmid Glycoprotein (AHSG).

Target: Alpha-2-Heremans Schmid Glycoprotein (AHSG)

Clonality: Polyclonal

Reactivity: Cow

Tested Applications: WB, IHC, IF/ICC

Host: Rabbit

Recommended dilutions: WB: 0.5-2 μg/ml, IHC: 5-20 μg/ml, IF/ICC: 5-20 μg/ml. Optimal dilutions/concentrations should be

determined by the end user.

Conjugation: Unconjugated

Immunogen: <u>abx652476</u> - Recombinant AHSG (Ile19-Ile359) expressed in E. coli

Form: Liquid

Purification: Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Buffer: PBS, pH 7.4, containing 0.02% NaN3, 50% glycerol.

Specificity: Cross-reacts with Human AHSG.

Concentration: 0.85 mg/ml

Note: This product is for research use only.

2 of 2

Abbexa LTD, Cambridge, UK · Phone: +44 (0) 1223 755950 · Fax: +44 (0) 1223 755951