Human Protein O-GlcNAcase / MGEA5 (OGA) Protein

Catalogue No.:abx068189



SDS-PAGE analysis of recombinant Human MGEA5 Protein.



Gene sequencing extract of recombinant Human MGEA5 Protein.

Human Meningioma Expressed Antigen 5 (MGEA5) is a recombinant Human protein produced in a Prokaryotic expression system (E. coli).

Target:	Protein O-GlcNAcase / MGEA5 (OGA)
Origin:	Human
Expression:	Recombinant
Tested Applications:	WB, SDS-PAGE
Host:	E. coli
Conjugation:	Unconjugated
Form:	Lyophilized
Purity:	> 90%
Reconstitution:	To keep the original salt concentration, we recommend reconstituting to the original concentration prior to lyophilization (see Concentration) in ddH_2O. If a lower concentration is required, dilute in 20 mM Tris, 150 mM NaCl, pH 8.0. If a higher concentration is required, the product can be reconstituted directly in 20 mM Tris, 150 mM NaCl, pH 8.0, though please note that this will change the overall salt concentration. The stock concentration should be between 0.1-1.0 mg/ml. Do not vortex.

Abbexa LTD, Cambridge, UK · Phone: +44 (0) 1223 755950 · Fax: +44 (0) 1223 755951
Abbexa LLC, Houston, TX USA · Phone: +1 832 327 7413
Abbexa BV, Leiden, NL
Website: www.abbexa.com · Email: info@abbexa.com

Storage:	Store at 2-8 °C for up to one month. Store at -80 °C for up to one year. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	O60502 (UniProt, ExPASy)
KEGG:	hsa:10724
String:	9606.ENSP00000354850
Molecular Weight:	Calculated MW: 27.5 kDa Observed MW (SDS-PAGE): 33 kDa Possible reasons why the actual band size differs from the predicted band size: 1. Splice variants. Alternative splicing may create different sized proteins from the same gene. 2. Relative charge. The composition of amino acids may affect the charge of the protein. 3. Post-translational modification. Phosphorylation, glycoslyation, methylation etc. may affect the band size. 4. Post-translational cleavage. Many proteins are synthesised as pro-proteins, and then cleaved to give the active form. 5. Polymerisation of the target protein. Dimerisation, multimerisation etc. will increase the band size observed.
Sequence Fragment:	Ser6-Lys240
Sequence:	SQATL EERESELSSN PAASAGASLE PPAAPAPGED NPAGAGGAAV AGAAGGARRF LCGVVEGFYG RPW VMEQRKE LFRRLQKWEL NTYLYAPKDD YKHRMFWREM YSVEEAEQLM TLISAAREYE IEFIYAISPG L DITFSNPKE VSTLKRKLDQ VSQFGCRSFA LLFDDIDHNM CAADKEVFSS FAHAQVSITN EIYQYLGEPE TFLFCPTEYC GTFCYPNVSQ SPYLRTVGEK
Tag:	N-terminal His tag
Buffer:	Prior to lyophilization: 20 mM Tris, 150 mM NaCl, pH 8.0, containing 0.01% Sarcosyl, 5% Trehalose.
Activity:	Not tested
Concentration:	Prior to lyophilization: 150 µg/ml
Note:	This product is for research use only. Not for human consumption, cosmetic, therapeutic or diagnostic use.

Abbexa LTD, Cambridge, UK · Phone: +44 (0) 1223 755950 · Fax: +44 (0) 1223 755951
Abbexa LLC, Houston, TX USA · Phone: +1 832 327 7413
Abbexa BV, Leiden, NL
Website: www.abbexa.com · Email: info@abbexa.com