



Activating Transcription Factor 4 (ATF4) Antibody


Catalogue No.: abx103393




Western blot analysis of recombinant Human ATF4.




IHC-P analysis of Human Prostate Gland Cancer Tissue, with DAB staining.



IHC-P analysis of Human Breast Cancer Tissue, with DAB staining.



IHC-P analysis of Human Stomach Cancer Tissue, with DAB staining.



IHC-P analysis of Human Skin Cancer Tissue, with DAB staining.

Datasheet

Version: 3.0.0
Revision date: 23 Feb 2025



Polyclonal Antibody to ATF4 (ATF4).

Target:	Activating Transcription Factor 4 (ATF4)
Clonality:	Polyclonal
Reactivity:	Human
Tested Applications:	WB, IHC, IF/ICC
Host:	Rabbit
Recommended dilutions:	WB: 0.01-2 µg/ml, IHC: 5-20 µg/ml, IF/ICC: 5-20 µg/ml. Optimal dilutions/concentrations should be determined by the end user.
Conjugation:	Unconjugated
Immunogen:	ATF4 (Thr130-Lys347)
Form:	Liquid
Purification:	Purified by antigen-specific affinity chromatography, followed by Protein A affinity chromatography.
Storage:	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
UniProt Primary AC:	P18848 (UniProt , ExpASY)
KEGG:	hsa:468
String:	9606.ENSP00000336790
Buffer:	0.01 M PBS, pH 7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration:	1 mg/ml
Note:	This product is for research use only.