



## Goat Anti-Rabbit IgG H&L Antibody (TRITC)


Catalogue No.: abx005567



Immunofluorescence analysis of HeLa cells, using Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) as the primary antibody at dilution of 1/100. The cells were incubated with the primary antibody overnight at 4°C. Secondary antibody: Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) at 1/200 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells, using Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) as the primary antibody at dilution of 1/100. The cells were incubated with the primary antibody overnight at 4°C. Secondary antibody: Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) at 1/100 dilution. Blue: DAPI for nuclear staining.



Flow cytometry:  $1 \times 10^6$  K-562 cells (negative control, left) and A-431 cells (right) were surface-stained with Purified Rabbit anti-Human E-Cadherin antibody (5  $\mu$ l/Test, orange line) or secondary antibody only (blue line). Non-fluorescently stained K-562 and A-431 cells were used as blank control (red line). Rhodamine (TRITC) Goat Anti-Rabbit IgG (H+L) (1/200) was used as a secondary antibody.

Goat Anti-Rabbit IgG (H+L) Antibody (TRITC) is a Goat antibody against Rabbit IgG (H+L). This product is for research use only, not for diagnostic or therapeutic use.

**Target:** Rabbit IgG H&L

**Clonality:** Polyclonal

**Reactivity:** Rabbit

**Tested Applications:** IHC, IF/ICC, FCM

**Host:** Goat

**Recommended dilutions:** IHC-P: 1/50 - 1/200, IF/ICC: 1/50 - 1/200, FCM: 1/50 - 1/200. Not tested in IHC-F. Optimal dilutions/concentrations should be determined by the end user.

# Datasheet

Version: 5.0.0  
Revision date: 30 Mar 2025



<b>Conjugation:</b>	TRITC
<b>Excitation/Emission:</b>	552/576
<b>Laser Line:</b>	550
<b>Immunogen:</b>	Rabbit IgG
<b>Isotype:</b>	IgG
<b>Form:</b>	Liquid
<b>Purification:</b>	Purified by affinity chromatography.
<b>Storage:</b>	Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.
<b>Buffer:</b>	PBS, pH 7.3, containing 0.025% sodium azide, 0.75% BSA, 50% glycerol.
<b>Concentration:</b>	> 0.2 mg/ml
<b>Note:</b>	THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION.

For Reference Only