

## Ubiquitin C-Terminal Hydrolase UCH37 (UCH37) Antibody

Catalogue No.: abx122753

Rabbit Polyclonal against the UCH37 protein.

|                               |  |
|-------------------------------|--|
| <b>Target:</b>                | Ubiquitin C-Terminal Hydrolase UCH37 (UCH37)   |
| <b>Clonality:</b>             | Polyclonal   |
| <b>Reactivity:</b>            | Human  |
| <b>Tested Applications:</b>   | ELISA, WB, IHC   |
| <b>Host:</b>                  | Rabbit   |
| <b>Recommended dilutions:</b> | ELISA: 1/20000 - 1/80000, WB: 1/500 - 1/2000, IHC: 1/100 - 1/200. Optimal dilutions/concentrations should be determined by the end user.   |
| <b>Conjugation:</b>           | Unconjugated   |
| <b>Immunogen:</b>             | A peptide corresponding to the C terminal region of human UCH37 protein.   |
| <b>Isotype:</b>               | IgG  |
| <b>Form:</b>                  | Lyophilized  |
| <b>Purification:</b>          | Purified by antigen affinity column chromatography.  |
| <b>Reconstitution:</b>        | Reconstitute in 100 µl of sterile distilled H <sub>2</sub> O with 50% glycerol.  |
| <b>Storage:</b>               | Store at -20 °C. Avoid repeated freeze/thaw cycles.  |
| <b>NCBI Accession:</b>        | NP_057068  |
| <b>Molecular Weight:</b>      | Observed MW: 36 kDa  |
| <b>Buffer:</b>                | Prior to lyophilization: 1% BSA and 0.02% NaN <sub>3</sub> .   |
| <b>Concentration:</b>         | Lyophilized form: Not applicable.<br>After reconstitution: 1 mg/ml.  |
| <b>Note:</b>                  | THIS PRODUCT IS FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC, THERAPEUTIC OR COSMETIC PROCEDURES. NOT FOR HUMAN OR ANIMAL CONSUMPTION. |