
Datasheet

Version: 2.0.0 Revision date: 18 Oct 2024



Androgen Receptor (AR) Antibody

Catalogue No.:abx011629



Androgen Receptor (AR) Antibody is a Rabbit Polyclonal antibody for the detection of androgen receptor.

The androgen receptor gene is more than 90 kb long and codes for a protein that has 3 major functional domains: the N-terminal domain, DNA-binding domain, and androgen-binding domain. The protein functions as a steroid-hormone activated transcription factor. Upon binding the hormone ligand, the receptor dissociates from accessory proteins, translocates into the nucleus, dimerizes, and then stimulates transcription of androgen responsive genes. This gene contains 2 polymorphic trinucleotide repeat segments that encode polyglutamine and polyglycine tracts in the N-terminal transactivation domain of its protein. Expansion of the polyglutamine tract from the normal 9-34 repeats to the pathogenic 38-62 repeats causes spinal bulbar muscular atrophy (SBMA, also known as Kennedy's disease). Mutations in this gene are also associated with complete androgen insensitivity (CAIS). Alternative splicing results in multiple transcript variants encoding different isoforms.

Target: Androgen Receptor (AR)

Clonality: Polyclonal

Reactivity: Human, Rat

Tested Applications: ELISA, WB, IHC

Host: Rabbit

Recommended dilutions: WB: 1/500 - 1/2000, IHC: 1/500 - 1/200. Optimal dilutions/concentrations should be determined by

the end user.

Conjugation: Unconjugated

Immunogen: androgen receptor

Isotype: IgG

Form: Liquid

Purity: ≥ 95% (SDS-PAGE)

Datasheet

Version: 2.0.0 Revision date: 18 Oct 2024



Purification: Purified by immunogen affinity chromatography.

Storage: Aliquot and store at -20°C. Avoid repeated freeze/thaw cycles.

Validity: 12 months.

UniProt Primary AC: P10275 (UniProt, ExPASy)

Gene Symbol: AR

GeneID: 367

OMIM: <u>300068</u>

HGNC: 644

KEGG: hsa:367

Ensembl: ENSG00000169083

String: <u>9606.ENSP00000363822</u>

Molecular Weight: Observed MW: 45 kDa, 75-80 kDa

Buffer: PBS, pH 7.3, with 0.02% sodium azide and 50% glycerol.

Concentration: 2 mg/ml

Note: This product is for research use only.